

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p>				
PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.				
1. REPORT DATE (DD-MM-YYYY) 10/19/2017	2. REPORT TYPE Poster	3. DATES COVERED (From - To) 10/19/2017-10/23/2017		
4. TITLE AND SUBTITLE Mediastinal Masses in Pediatric Patient with Ewing's Sarcoma		5a. CONTRACT NUMBER 5b. GRANT NUMBER 5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Maj Michael F D'Amore		5d. PROJECT NUMBER 5e. TASK NUMBER 5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg 4430 JBSA-Lackland, TX 78236-9908 210-292-7141			8. PERFORMING ORGANIZATION REPORT NUMBER 17410	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg 4430 JBSA-Lackland, TX 78236-9908 210-292-7141			10. SPONSOR/MONITOR'S ACRONYM(S) 11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release. Distribution is unlimited.				
13. SUPPLEMENTARY NOTES American Society of Anesthesiologists Annual Meeting 2017 (ASA), Boston MA 19-23 Oct 2017				
14. ABSTRACT				
15. SUBJECT TERMS				
16. SECURITY CLASSIFICATION OF: a. REPORT b. ABSTRACT c. THIS PAGE		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Claricel Longoria 19b. TELEPHONE NUMBER (Include area code) 210-292-7141



Mediastinal Masses in Pediatric Patient with Ewing's Sarcoma

Michael D'Amore M.D., Maj, USAF, MC, FS; Angela McElrath M.D., Maj, USAF, MC, FS; Christina Rhodes D.O.
San Antonio Uniformed Services Health Education Consortium (SAUSHEC) Anesthesiology Program



Case Description

SyOm ASA 4 with history of Ewing's sarcoma involving T1-L1 now 18mo off chemotherapy presenting with multiple large mediastinal hilar, and pleural masses concerning for metastasis found on surveillance imaging. CT scan revealed mass effect (compression & shift) on the trachea and left pulmonary vein secondary to 6x4x3cm para-tracheal mass. Additional mediastinal masses were noted and associated with right bronchial compression and large right pleural effusion.

Preoperative Course

PreOp Vitals: 106/46, HR 103, RR 20, SpO₂ 100%. TTE accomplished and revealed normal biventricular systolic function, no outflow tract obstruction. Peripheral IVs x2 placed by PICC team the night prior. OR prepped with rigid bronchoscopy tray and general surgeon in room at induction in the event an emergent rigid bronchoscopy is required.

Anesthetic Management

Pt taken to OR for inhalational induction and LMA placement. Pt maintained on sevoflurane while breathing spontaneously. Pt taken to MRI for scan. After MRI the pt returned back to OR still under general anesthesia, breathing spontaneously. Additional peripheral IV was placed and pre-procedure arterial line placed for hemodynamic monitoring. Laryngoscopy was performed and ETT 5.5 placed to secure airway with surgeons present and rigid bronchoscopy tray available. Spontaneous ventilation maintained throughout and general anesthesia maintained with propofol infusion, dexamethasone, ketamine, and volatile anesthetics. Anesthesiologist vigilance and preparation prevented potential airway obstruction and cardiovascular collapse.


Mediastinal Mass Obstruction Physiology

- 4 compartments of Mediastinum: Anterior, Middle (visceral), Posterior (Paravertebral), Superior
- Most common masses in the anterior mediastinum are the four "T's"
 - Thymoma, teratoma, Thyroid carcinoma and Terribile lymphoma.
- Anatomic Effects of Mediastinal Mass
 - Compression of cardiac structures- rhythm disturbances, distal flow reduction
 - Compression of vascular structures- compression of SVC or distal flow reduction
 - Compression and collapse of airway structures
- Lung volumes reduced to 500-1500cc- Decrease in tracheobronchial diameter
- Bronchial smooth muscle relaxation for easier compressibility
- Compression eliminates the normal trans-pulmonary pressure gradient causing enhanced extrinsic compression and decreased airway diameter.

Pitfalls on Anesthetic Induction

- Asymptomatic patient- no symptoms
- Mid- Supine with some cough/pressure sensation
- Severe- Cannot tolerate supine

CT scan and determination of symptoms- patients are deemed safe, unsafe, or uncertain for anesthesia as follows:



- Safe**
 - Asymptomatic adult, CT < 50% tracheobronchial obstruction
 - Severely symptomatic adult or child
 - Children with CT tracheobronchial obstruction > 50%, regardless of symptoms
- Unsafe**
 - Mild- Supine with some cough/pressure sensation
 - Moderate- Supine for short periods
 - Severe- Cannot tolerate supine

Pediatric vs Adult

- Emergency Response to Airway Compression**
 - Anesthetic deaths mainly reported in children
 - More compressible cartilaginous structure of the airway
 - Underestimation of severity of symptoms due to obtaining a clear history
 - Aware fiberoptic intubation to obtain distal airway is not an option in most children
 - Children with 50% or greater tracheo-bronchial compression have a greater likelihood of severe complications with general anesthesia
- Mediastinal Mass Operative Management**
 - Cannulate femoral vessels for possible need for Cardio-Pulmonary Bypass in unsafe and uncertain category
 - Adjust patient position- elevated head of bed and partial or full right lateral decubitus position to maintain airway patency and reduce cardiac and vascular compression
 - Invasive arterial monitoring- potential cardiorespiratory instability
 - Maintain spontaneous ventilation if possible
 - Awake fiber optic intubation
 - Inhalational induction
 - Secure airway beyond stenosis if possible
 - CPAP may help with maintaining functional residual capacity
 - Short acting anesthetics and paralytics (avoid if able)
 - Rigid bronchoscopy and surgeon available at induction

References:

- Miller's Anesthesia, 7th ed. Philadelphia: Elsevier Saunders; 2014. 1997-1998. Anesthetic considerations for patients with anterior mediastinal masses.
- Anesthesiology Clin. 2008 Jun; 26 (2): 305-14.
- Surprenant P, Pernow B. Anesthesia and Physiology in Anesthetic Practice. St. Louis: 2015. Chapter 24.
- Arteriovenous Malformations: Anesthetic Considerations in Mediastinal Mass during Thoracic Department of Cardio-Anesthesiology (GIVE) & R. Untalan. Nov 2014. CTVA-2014.
- M. H. Cho, X. J. Lin and S. Goh. Anterior mediastinal masses: an anaesthetic challenge. Anesthesia. 1999, 54, 697-705.
- Singh, Peter, Kausik. (2007). Management of the patient with a large anterior mediastinal mass: recurring myths. Current Opinion in Anesthesiology. 2001; 12: 3-10.
- Halmér, György. (2007). Anesthetic management for the child with a mediastinal mass. Pediatric Anesthesia. 07/2006; Vol 14: 55-57.
- Pullicino, J. Anesthesia for patients with mediastinal masses. Canadian Journal of Anesthesia. 11/1/1989; Vol 36(6): 681-686.

Disclaimer: The views(s) expressed herein are those of the author(s) and do not reflect the official policy or position of San Antonio Military Medical Center, the U.S. Army Medical Department, the U.S. Army Office of the Surgeon General, the Department of the Air Force, the Department of the Army or the Department of Defense or the U.S. Government.